arXiv:1208.6228v2 [math.PR] 17 Sep 2012

The distribution function of the distance between two random points in a right-angled triangle

Uwe Bäsel

Abstract

In this paper we obtain the density function and the distribution function of the distance between two uniformly and independently distributed random points in any right-angled triangle. The density function is derived from the chord length distribution function using Piefke's formula. We conclude results for random distances between two congruent right-angled triangles together forming a rectangle.

2010 Mathematics Subject Classification: 60D05, 52A22

Keywords: Geometric probability, random sets, integral geometry, chord length distribution function, random distances, distance distribution function, right-angled triangles, right triangle, Piefke formula

1 Introduction

A random line g intersecting a convex set $\mathcal{K}$ in the plane produces a chord of $\mathcal{K}$. The length s of this chord is a random variable. The chord length distribution function of a regular triangle was calculated by Sulanke [14, p. 57]. Harutyunyan and Ohanyan [8] calculated the chord length distribution function for regular polygons (see also Bäsel [2, pp. 2-9]).

The distance t between two points chosen independently and uniformly at random from $\mathcal{K}$ is also a random variable. Borel [3] considered random distances in elementary geometric figures such as triangles, squares and so on (see [11, p. 163]). The expectations of the distance between two random points for an equilateral triangle and a rectangle are to be found in [13, p. 49]. Ghosh [6] derived the distance density function for a rectangle. Bäsel [2, pp. 10-16] derived the density function and the distribution function of the distance for regular polygons from its chord length distribution function using Piefke's formula [12, p. 130]. There are a lot of results concerning the distance t within a convex set or in two convex sets, see Chapter 2 in [10].

For practical applications in physics, material sciences, operations research, communication networks and other fields see [7], [9] and [15].

The first aim of the present paper is to derive the chord length distribution function for any right-angled triangle. The second aim is to conclude the distribution function of the distance between two points chosen independently and uniformly from a right-angled triangle.

$A = (0, b)$, $B = (a, 0)$ and $C = (0, 0)$ are the vertices of the right-angled triangle (see Fig. 1b). We assume $|\overline{AC}| = b \geq a = |\overline{BC}|$ and denote this triangle by $\mathcal{T}_{a,b}$. Clearly, it is possible to bring every right-angled triangle in such a position by use of Euclidean motions (translations and rotations) and, if necessary, a reflection. These operations do not influence the chord length distribution and the point distance distribution. $c = |\overline{AB}|$ is the length of the hypotenuse and $\alpha = \arctan(a/b)$ the inner angle between $\overline{AB}$ and $\overline{AC}$. $u = a + b + c$ is the length of the perimeter of $\mathcal{T}_{a,b}$, $\tilde{A} = ab/2$ its area and $h = a \cos \alpha$ the height over $\overline{AB}$. The hypotenuse $\overline{AB}$ of the triangle $\mathcal{T}_{a,b}$ is the diameter of its circumcircle (converse of Thales' theorem).

2 Chord length distribution function

A straight line g in the plane is determined by the angle ψ, $0 \leq \psi < 2\pi$, that the direction perpendicular to g makes with a fixed direction (e.g. the x-axis) and by its distance p, $0 \leq p < \infty$, from the origin O:

$$g = g(p, \psi) = \{(x, y) \in \mathbb{R}^2 : \, x \cos \psi + y \sin \psi = p\} \, .$$

The measure μ of a set of lines $g(p, \psi)$ is defined by the integral, over the set, of the differential form $\mathrm{d}g = \mathrm{d}p \, \mathrm{d}\psi$. Up to a constant factor, this measure is the only one that is invariant under motions in the Euclidean plane [13, p. 28].

We use the chord length distribution function of $\mathcal{T}_{a,b}$ in the form

$$F(s) = \frac{1}{u} \, \mu(\{g : \, g \cap \mathcal{T}_{a,b} \neq \emptyset, \, |\chi(g)| \leq s\}) \, ,$$

where $\chi(g) = g \cap \mathcal{T}_{a,b}$ is the chord of $\mathcal{T}_{a,b}$, produced by the line g, and $|\chi(g)|$ the length of $\chi(g)$ (see [5, p. 863] and [1, p. 161]). (The measure of all lines g that intersect a convex set is equal to its perimeter [13, p. 30].) So it remains to calculate the measure of all lines that produce a chord of length $|\chi(g)| \leq s$. Using the abbreviation

$$\mathcal{S}(g, s) := \{g : \, g \cap \mathcal{T}_{a,b} \neq \emptyset, \, |\chi(g)| \leq s\} \, ,$$

we have

$$\mu(\mathcal{S}(g, s)) = \int_{\mathcal{S}(g,s)} \mathrm{d}g = \int_{\mathcal{S}(g,s)} \mathrm{d}p \, \mathrm{d}\psi \, .$$

Instead of the angle ψ it is possible to use the angle ϕ that a line makes with the positive x-axis, hence

$$\mu(\mathcal{S}(g, s)) = \int_{\mathcal{S}(g,s)} \mathrm{d}p \, \mathrm{d}\phi \, .$$

Theorem 1. *The chord length distribution function F of $\mathcal{T}_{a,b}$ is given by*

$$F(s) = \begin{cases} 0 & \text{if } \ s < 0\,, \\ H_1(s)/u & \text{if } \ 0 \le s < h\,, \\ H_2(s)/u & \text{if } \ h \le s < a\,, \\ H_3(s)/u & \text{if } \ a \le s < b\,, \\ H_4(s)/u & \text{if } \ b \le s < c\,, \\ 1 & \text{if } \ s \ge c\,, \end{cases}$$

where

$$\begin{aligned} H_1(s) &= \frac{\Theta_1\, s}{4}\,, \\ H_2(s) &= \frac{\Theta_2\, s}{4} + \Theta_3\, L_1(s,h) + c\, L_2(s,h)\,, \\ H_3(s) &= a + \Theta_4\, s + \frac{\Theta_3}{2}\, L_1(s,h) + \frac{c}{2}\, L_2(s,h) + \frac{b}{2a}\, L_1(s,a) + \frac{b}{2}\, L_2(s,a)\,, \\ H_4(s) &= \Theta_5 + \frac{\Theta_6\, s}{4} + \frac{b}{2a}\, L_1(s,a) + \frac{b}{2}\, L_2(s,a) + \frac{a}{2b}\, L_1(s,b) + \frac{a}{2}\, L_2(s,b) \end{aligned}$$

with

$$\begin{aligned} &\Theta_1 = \frac{a}{b}\,(2\alpha + \pi) + \frac{2b}{a}\,(\pi - \alpha) + 6\ , \quad \Theta_2 = \frac{a}{b}\,(2\alpha - \pi) - \frac{2b}{a}\,\alpha + 6\,, \\ &\Theta_3 = \frac{a}{b} + \frac{b}{a}\ , \quad \Theta_4 = 1 - \frac{b}{a}\,\alpha\ , \quad \Theta_5 = a + b\ , \quad \Theta_6 = \frac{a}{b}\,(2\alpha - \pi) - \frac{2b}{a}\,\alpha + 2 \end{aligned}$$

and

$$L_1(s,a) = s \arcsin\frac{a}{s}\ , \quad L_2(s,a) = \sqrt{1 - \left(\frac{a}{s}\right)^2}\ .$$

Proof. In the following we need the width w of $\mathcal{T}_{a,b}$ in the direction perpendicular to the angle ϕ that is given by:

$$w(\phi) = \begin{cases} w_1(\phi) & \text{if} \quad 0 \le \phi < \pi/2\,, \\ w_2(\phi) & \text{if} \quad \pi/2 \le \phi < \pi/2 + \alpha\,, \\ w_3(\phi) & \text{if} \quad \pi/2 + \alpha \le \phi \le \pi\,, \end{cases}$$

where

$$w_1(\phi) = a\,\frac{\cos(\phi - \alpha)}{\sin\alpha}\ , \quad w_2(\phi) = a \sin\phi\ , \quad w_3(\phi) = -a\,\frac{\cos\phi}{\tan\alpha}\ .$$

We have to distinguish the cases $0 \le s < h$, $h \le s < a$, $a \le s < b$ and $b \le s < c$, and put

$$\begin{aligned} &H_1(s) = \mu(\mathcal{S}(g,s)) \ \text{ if } \ 0 \le s < h\ , \quad H_2(s) = \mu(\mathcal{S}(g,s)) \ \text{ if } \ h \le s < a\ , \\ &H_3(s) = \mu(\mathcal{S}(g,s)) \ \text{ if } \ a \le s < b\ , \quad H_4(s) = \mu(\mathcal{S}(g,s)) \ \text{ if } \ b \le s < c\ . \end{aligned}$$

We demonstrate the calculation of H_2. Here it is necessary to distinguish the following intervals of ϕ (see Fig. 1):

a) $0 \le \phi < \phi_1 = \alpha - \arccos\left((a/s)\cos\alpha\right)$: All lines g with fixed angle ϕ produce a chord χ of length $|\chi(g)| \le s$ if they are lying in one of the two strips of breadth

$$b_1(s,\phi) = \frac{s}{2}\,\frac{\sin(2\phi-\alpha)+\sin\alpha}{\cos\alpha} \quad \text{and} \quad b_2(s,\phi) = \frac{s}{2}\,\frac{\cos(2\phi-\alpha)+\cos\alpha}{\sin\alpha}$$

respectively.

b) $\phi_1 \le \phi < \phi_2 = \alpha + \arccos\left((a/s)\cos\alpha\right)$: All lines g with fixed angle ϕ and $g \cap \mathcal{T}_{a,b} \ne \emptyset$ produce a chord of length $|\chi(g)| \le s$. These lines are lying in a strip of breadth $w_1(\phi)$.

c) $\phi_2 \le \phi < \pi/2$: See $0 \le \phi < \phi_1$.

d) $\pi/2 \le \phi < \pi/2 + \alpha$: All lines g with fixed angle ϕ produce a chord of length $\le s$ if they are lying in one of the two strips of breadth $b_1(s,\phi)$ and

$$b_3(s,\phi) = -\frac{s}{2}\sin 2\phi$$

respectively.

e) $\pi/2 + \alpha \le \phi < \pi$: All lines g with fixed angle ϕ, lying in one of the two strips of breadth $b_2(s,\phi)$ and $b_3(s,\phi)$ repectively, produce a chord of length $\le s$.

Therefore, using the abbreviations b_i and w_i for $b_i(s,\phi)$ and $w_i(\phi)$ respectively,

$$\begin{aligned} H_2(s) = \int_{\mathcal{S}(g,s)} \mathrm{d}p\,\mathrm{d}\phi &= \int_0^{\phi_1} (b_1+b_2)\,\mathrm{d}\phi + \int_{\phi_1}^{\phi_2} w_1\,\mathrm{d}\phi + \int_{\phi_2}^{\pi/2} (b_1+b_2)\,\mathrm{d}\phi \\ &\quad + \int_{\pi/2}^{\pi/2+\alpha} (b_1+b_3)\,\mathrm{d}\phi + \int_{\pi/2+\alpha}^{\pi} (b_2+b_3)\,\mathrm{d}\phi\,. \end{aligned}$$

With the angles

$$\phi_3 = \pi - \arccos\frac{a}{s} \quad \text{and} \quad \phi_4 = \pi - \arcsin\left(\frac{a}{s}\cot\alpha\right)$$

we analogously get

$$\begin{aligned} H_1(s) &= \int_0^{\pi/2} (b_1+b_2)\,\mathrm{d}\phi + \int_{\pi/2}^{\pi/2+\alpha} (b_1+b_3)\,\mathrm{d}\phi + \int_{\pi/2+\alpha}^{\pi} (b_2+b_3)\,\mathrm{d}\phi\,, \\ H_3(s) &= \int_0^{\phi_2} w_1\,\mathrm{d}\phi + \int_{\phi_2}^{\pi/2} (b_1+b_2)\,\mathrm{d}\phi + \int_{\pi/2}^{\pi/2+\alpha} (b_1+b_3)\,\mathrm{d}\phi \\ &\quad + \int_{\pi/2+\alpha}^{\phi_3} (b_2+b_3)\,\mathrm{d}\phi + \int_{\phi_3}^{\pi} w_3\,\mathrm{d}\phi\,, \end{aligned}$$



Figure 1: Calculation of H_2

$$H_4(s) = \int_0^{\pi/2} w_1 \,\mathrm{d}\phi + \int_{\pi/2}^{\phi_4} w_2 \,\mathrm{d}\phi + \int_{\phi_4}^{\pi/2+\alpha} (b_1 + b_3) \,\mathrm{d}\phi$$
$$+ \int_{\pi/2+\alpha}^{\phi_3} (b_2 + b_3) \,\mathrm{d}\phi + \int_{\phi_3}^{\pi} w_3 \,\mathrm{d}\phi \,.$$

Using the indefinite integrals

$$\begin{aligned}
\int b_1(s,\phi)\,\mathrm{d}\phi &= \frac{s}{4}\,\frac{2\phi\sin\alpha - \cos(2\phi-\alpha)}{\cos\alpha}\,,\\
\int b_2(s,\phi)\,\mathrm{d}\phi &= \frac{s}{4}\,\frac{2\phi\cos\alpha + \sin(2\phi-\alpha)}{\sin\alpha}\,,\\
\int b_3(s,\phi)\,\mathrm{d}\phi &= \frac{s}{2}\,\cos^2\phi\,,\\
\int w_1(\phi)\,\mathrm{d}\phi \ &= \frac{a\sin(\phi-\alpha)}{\sin\alpha} = a(\cot\alpha\sin\phi - \cos\phi)\,,\\
\int w_2(\phi)\,\mathrm{d}\phi \ &= -\,a\cos\phi\,, \quad \int w_3(\phi)\,\mathrm{d}\phi = -\frac{a\sin\phi}{\tan\alpha}\,,
\end{aligned}$$

one finds the result of Theorem 1. ☐

Remark. It is easy to derive the chord length density function from Theorem 1.

3 Point distances

Theorem 2. *The density function g of the distance t between two random points in $\mathcal{T}_{a,b}$ is given by*

$$g(t) = \begin{cases} \dfrac{2t}{\tilde{A}}\left[\pi + \dfrac{1}{\tilde{A}}\,(J^*(t) - ut)\right] & \textit{if} \quad 0 \le t \le c\,,\\ 0 & \textit{else}\,, \end{cases}$$

where

$$J^*(t) = \begin{cases} J_1^*(0,t) & \textit{if} \quad 0 \le t < h\,,\\ J_1^*(0,h) + J_2^*(h,t) & \textit{if} \quad h \le t < a\,,\\ J_1^*(0,h) + J_2^*(h,a) + J_3^*(a,t) & \textit{if} \quad a \le t < b\,,\\ J_1^*(0,h) + J_2^*(h,a) + J_3^*(a,b) + J_4^*(b,t) & \textit{if} \quad b \le t \le c \end{cases}$$

with

$$J_k^*(s,t) = H_k^*(t) - H_k^*(s)\,,$$

where

$$
\begin{aligned}
H_1^*(t) &= \frac{\Theta_1\, t^2}{8}\,,\\
H_2^*(t) &= \frac{\Theta_2\, t^2}{8} + \Theta_3\, L_1^*(t,h) + c\, L_2^*(t,h)\,,\\
H_3^*(t) &= at + \frac{\Theta_4\, t^2}{2} + \frac{\Theta_3}{2}\, L_1^*(t,h) + \frac{c}{2}\, L_2^*(t,h) + \frac{b}{2a}\, L_1^*(t,a) + \frac{b}{2}\, L_2^*(t,a)\,,\\
H_4^*(t) &= \Theta_5\, t + \frac{\Theta_6\, t^2}{8} + \frac{b}{2a}\, L_1^*(t,a) + \frac{b}{2}\, L_2^*(t,a) + \frac{a}{2b}\, L_1^*(t,b) + \frac{a}{2}\, L_2^*(t,b)
\end{aligned}
$$

with

$$L_1^*(t,a) = \frac{1}{2}\left(a\sqrt{t^2-a^2} + t^2 \arcsin\frac{a}{t}\right)\,,\quad L_2^*(t,a) = \sqrt{t^2-a^2} + a \arcsin\frac{a}{t}\,.$$

Proof. According to a theorem of Piefke [12, p. 130], the density function of the distance between two random points is given by

$$g(t) = \frac{2ut}{\tilde{A}^2}\int_t^c (s-t) f(s)\,\mathrm{d}s\,,$$

where f is the density function of the chord length. Following [2, pp. 11/12], we obtain

$$g(t) = \frac{2t}{\tilde{A}}\left[\pi - \frac{u}{\tilde{A}}\left(t - \int_0^t F(s)\,\mathrm{d}s\right)\right] = \frac{2t}{\tilde{A}}\left[\pi + \frac{1}{\tilde{A}}\left(J^*(t) - ut\right)\right]$$

with

$$J^*(t) := u\int_0^t F(s)\,\mathrm{d}s\,.$$

This yields for $0 \le t < h$

$$J^*(t) = \int_0^t H_1(s)\,\mathrm{d}s = H_1^*(t) - H_1^*(0) = J_1^*(0,t)\,,$$

and for $h \le t < a$

$$
\begin{aligned}
J^*(t) = u\int_0^t F(s)\,\mathrm{d}s &= \int_0^h H_1(s)\,\mathrm{d}s + \int_h^t H_2(s)\,\mathrm{d}s\\
&= H_1^*(h) - H_1^*(0) + H_2^*(t) - H_2^*(h) = J_1^*(0,h) + J_2^*(h,t)\,.
\end{aligned}
$$

The calculations of $J^*(t)$ for the intervals $a \le t < b$ and $b \le t < c$ can be carried out in the same manner.

Now we determine the indefinite integrals of $L_1(t,a)$ and $L_2(t,a)$. Using integration by parts, we find

$$
\begin{aligned}
L_1^*(t,a) = \int L_1(t,a)\,\mathrm{d}t &= \int t \arcsin\frac{a}{t}\,\mathrm{d}t\\
&= \frac{1}{2}\left(t^2 \arcsin\frac{a}{t} + a\int \frac{t}{\sqrt{t^2-a^2}}\,\mathrm{d}t\right).
\end{aligned}
$$

Clearly,

$$\int \frac{t}{\sqrt{t^2-a^2}}\,\mathrm{d}t = \sqrt{t^2-a^2}$$

and therefore

$$L_1^*(t,a) = \frac{1}{2}\left(a\sqrt{t^2-a^2} + t^2 \arcsin\frac{a}{t}\right).$$

Since $t \geq a > 0$ in the present cases,

$$L_2^*(t,a) = \int L_2(t,a)\,\mathrm{d}t = \int \sqrt{1-\left(\frac{a}{t}\right)^2}\,\mathrm{d}t = \int \frac{\sqrt{t^2-a^2}}{t}\,\mathrm{d}t\,,$$

and (cf. [4, p. 48, Eq. 217])

$$L_2^*(t,a) = \sqrt{t^2-a^2} + a \arcsin\frac{a}{t}\,. \qquad \square$$

Corollary 1. *The distribution function G of the distance t between two random points in $\mathcal{T}_{a,b}$ is given by*

$$G(t) = \begin{cases} 0 & \textit{if} \quad t < 0\,, \\ \dfrac{1}{\tilde{A}}\left[t^2\left(\pi - \dfrac{2u}{3\tilde{A}}\,t\right) + \dfrac{2}{\tilde{A}}\,J^\natural(t)\right] & \textit{if} \quad 0 \leq t < c\,, \\ 1 & \textit{if} \quad t \geq c \end{cases}$$

with

$$J^\natural(t) = \begin{cases} K_1(t) & \textit{if} \quad 0 \leq t < h\,, \\ K_1(h) + K_2(t) & \textit{if} \quad h \leq t < a\,, \\ K_1(h) + K_2(a) + K_3(t) & \textit{if} \quad a \leq t < b\,, \\ K_1(h) + K_2(a) + K_3(b) + K_4(t) & \textit{if} \quad b \leq t < c\,, \end{cases}$$

where

$$K_1(t) = J_1^\natural(0,t)\,,$$

$$K_2(t) = \frac{1}{2}\left(t^2-h^2\right)\left[J_1^*(0,h) - H_2^*(h)\right] + J_2^\natural(h,t)\,,$$

$$K_3(t) = \frac{1}{2}\left(t^2-a^2\right)\left[J_1^*(0,h) + J_2^*(h,a) - H_3^*(a)\right] + J_3^\natural(a,t)\,,$$

$$K_4(t) = \frac{1}{2}\left(t^2-b^2\right)\left[J_1^*(0,h) + J_2^*(h,a) + J_3^*(a,b) - H_4^*(b)\right] + J_4^\natural(b,t)$$

with J_k^* and H_k^* according to Theorem 2 and

$$J_k^\natural(s,t) = H_k^\natural(t) - H_k^\natural(s)\,,$$

where

$$
\begin{aligned}
H_1^\natural(t) &= \frac{\Theta_1\, t^4}{32}\,,\\
H_2^\natural(t) &= \frac{\Theta_2\, t^4}{32} + \Theta_3\, L_1^\natural(t,h) + c\, L_2^\natural(t,h)\,,\\
H_3^\natural(t) &= \frac{at^3}{3} + \frac{\Theta_4\, t^4}{8} + \frac{\Theta_3}{2}\, L_1^\natural(t,h) + \frac{c}{2}\, L_2^\natural(t,h) + \frac{b}{2a}\, L_1^\natural(t,a) + \frac{b}{2}\, L_2^\natural(t,a)\,,\\
H_4^\natural(t) &= \frac{\Theta_5\, t^3}{3} + \frac{\Theta_6\, t^4}{32} + \frac{b}{2a}\, L_1^\natural(t,a) + \frac{b}{2}\, L_2^\natural(t,a) + \frac{a}{2b}\, L_1^\natural(t,b) + \frac{a}{2}\, L_2^\natural(t,b)
\end{aligned}
$$

with

$$
\begin{aligned}
L_1^\natural(t,a) &= \frac{1}{8}\left[\frac{5a}{3}\left(t^2-a^2\right)^{3/2} + a^3\sqrt{t^2-a^2} + t^4 \arcsin\frac{a}{t}\right]\,,\\
L_2^\natural(t,a) &= \frac{1}{3}\left(t^2-a^2\right)^{3/2} + \frac{a}{2}\left(a\sqrt{t^2-a^2} + t^2\,\arcsin\frac{a}{t}\right)\,.
\end{aligned}
$$

Proof. For $0 \le t < c$ one gets

$$
\begin{aligned}
G(t) &= \int_0^t g(\tau)\,\mathrm{d}\tau = \int_0^t \left(\frac{2\pi\tau}{\tilde{A}} - \frac{2u\tau^2}{\tilde{A}^2} + \frac{2u\tau}{\tilde{A}^2}\int_0^\tau F(s)\,\mathrm{d}s\right)\mathrm{d}\tau\\
&= \frac{\pi t^2}{\tilde{A}} - \frac{2ut^3}{3\tilde{A}^2} + \frac{2u}{\tilde{A}^2}\int_0^t \tau\left(\int_0^\tau F(s)\,\mathrm{d}s\right)\mathrm{d}\tau\\
&= \frac{\pi t^2}{\tilde{A}} - \frac{2ut^3}{3\tilde{A}^2} + \frac{2}{\tilde{A}^2}\int_0^t \tau J^*(\tau)\,\mathrm{d}\tau = \frac{1}{\tilde{A}}\left[t^2\left(\pi - \frac{2u}{3\tilde{A}}\,t\right) + \frac{2}{\tilde{A}}\,J^\natural(t)\right]
\end{aligned}
$$

with

$$
J^\natural(t) := \int_0^t sJ^*(s)\,\mathrm{d}s\,.
$$

It remains to calculate $J^\natural(t)$. For $0 \le t < h$ we have

$$
\begin{aligned}
J^\natural(t) &= \int_0^t sJ_1^*(0,s)\,\mathrm{d}s = \int_0^t s\left[H_1^*(s) - H_1^*(0)\right]\mathrm{d}s\\
&= H_1^\natural(t) - H_1^\natural(0) - \frac{1}{2}\,t^2 H_1^*(0) = H_1^\natural(t) - H_1^\natural(0) - 0 = J_1^\natural(0,t) = K_1(t)\,.
\end{aligned}
$$

For $h \le t < a$ we find

$$
\begin{aligned}
J^\natural(t) &= K_1(h) + \int_h^t s\left[J_1^*(0,h) + J_2^*(h,s)\right]\mathrm{d}s\\
&= K_1(h) + \frac{1}{2}\left(t^2-h^2\right)J_1^*(0,h) + \int_h^t sJ_2^*(h,s)\,\mathrm{d}s\,.
\end{aligned}
$$

With

$$
\int_h^t sJ_2^*(h,s)\,\mathrm{d}s = \int_h^t s\left[H_2^*(s) - H_2^*(h)\right]\mathrm{d}s = \int_h^t sH_2^*(s)\,\mathrm{d}s - H_2^*(h)\int_h^t s\,\mathrm{d}s
$$

$$
\begin{aligned}
&= H_2^\natural(t) - H_2^\natural(h) - \frac{1}{2}\left(t^2 - h^2\right) H_2^*(h) \\
&= J_2^\natural(h,t) - \frac{1}{2}\left(t^2 - h^2\right) H_2^*(h)
\end{aligned}
$$

it follows that

$$
J^\natural(t) = K_1(h) + \frac{1}{2}\left(t^2 - h^2\right)\left[J_1^*(0,h) - H_2^*(h)\right] + J_2^\natural(h,t) = K_1(h) + K_2(t)\,.
$$

The calculation applies analogously to the intervals $a \le t < b$ and $b \le t < c$.

Now we calculate the indefinite integrals of $L_1^*(t,a)$ and $L_2^*(t,a)$. We have

$$
\begin{aligned}
L_1^\natural(t,a) &= \int t L_1^*(t,a)\,\mathrm{d}t = \frac{1}{2}\int t\left(a\sqrt{t^2 - a^2} + t^2 \arcsin\frac{a}{t}\right)\mathrm{d}t \\
&= \frac{1}{2}\left(a\int t\,\sqrt{t^2 - a^2}\,\mathrm{d}t + \int t^3 \arcsin\frac{a}{t}\,\mathrm{d}t\right)
\end{aligned}
$$

with

$$
\int t\,\sqrt{t^2 - a^2}\,\mathrm{d}t = \frac{1}{3}\left(t^2 - a^2\right)^{3/2} \quad [4, \text{p. 47, Eq. 214}]\,.
$$

Using integration by parts, we find

$$
\int t^3 \arcsin\frac{a}{t}\,\mathrm{d}t = \frac{1}{4}\left(t^4 \arcsin\frac{a}{t} + a\int \frac{t^2}{\sqrt{1 - (a/t)^2}}\,\mathrm{d}t\right).
$$

Since $t \ge a > 0$ in the present cases,

$$
\int \frac{t^2}{\sqrt{1 - (a/t)^2}}\,\mathrm{d}t = \int \frac{t^3}{\sqrt{t^2 - a^2}}\,\mathrm{d}t\,,
$$

and

$$
\int \frac{t^3}{\sqrt{t^2 - a^2}}\,\mathrm{d}t = \frac{1}{3}\left(t^2 - a^2\right)^{3/2} + a^2\,\sqrt{t^2 - a^2} \quad [4, \text{p. 48, Eq. 223}]\,.
$$

This yields

$$
L_1^\natural(t,a) = \frac{1}{8}\left[\frac{5a}{3}\left(t^2 - a^2\right)^{3/2} + a^3\,\sqrt{t^2 - a^2} + t^4 \arcsin\frac{a}{t}\right].
$$

Further,

$$
\begin{aligned}
L_2^\natural(t,a) &= \int t L_2^*(t,a)\,\mathrm{d}t = \int t\sqrt{t^2 - a^2}\,\mathrm{d}t + a\int t \arcsin\frac{a}{t}\,\mathrm{d}t \\
&= \frac{1}{3}\left(t^2 - a^2\right)^{3/2} + \frac{a}{2}\left(a\sqrt{t^2 - a^2} + t^2\,\arcsin\frac{a}{t}\right)
\end{aligned}
$$

(see the calculations of $L_1^\natural(t,a)$ and $L_1^*(t,a)$). $\square$

4 Two triangles

Now considering a rectangle $\mathcal{R}_{a,b}$ of side lengths a and b ($a \le b$) as made up of two congruent copies of the triangle $\mathcal{T}_{a,b}$ with different orientation and sharing their hypotenuses. Let g_2 and G_2 denote respectively the density function and the distribution function of the distance between two uniformly and independently distributed points P_1 and P_2, one in each of the triangles.

Using the *indirect method* (see Ghosh [6, pp. 22/23]), it is possible to calculate g_2 from the density function $g_{\mathcal{R}}$ of the distance between two random points inside $\mathcal{R}_{a,b}$ and the density function g of $\mathcal{T}_{a,b}$ (according to Theorem 2). The probability of P_1 and P_2 belonging to the same triangle is equal to the probability of P_1 and P_2 belonging to different triangles, each probability being 1/2. Therefore,

$$g_{\mathcal{R}} = \frac{1}{2}\,(g + g_2)\,, \quad \text{whence} \quad g_2 = 2\,g_{\mathcal{R}} - g \quad \text{and} \quad G_2 = 2\,G_{\mathcal{R}} - G\,,$$

where $G_{\mathcal{R}}$ denotes the corresponding distribution function of $g_{\mathcal{R}}$. As already mentioned, $g_{\mathcal{R}}$ was found by Ghosh [6] (Theorem 2, pp. 18/19). Denoting by $A_{\mathcal{R}} = ab$ the area of $\mathcal{R}_{a,b}$ and by $u_{\mathcal{R}} = 2(a+b)$ its perimeter, we can write Ghosh's result as

$$g_{\mathcal{R}}(t) = \begin{cases} \dfrac{2t}{A_{\mathcal{R}}^2}\, J_{\mathcal{R}}^*(t) & \text{if} \quad 0 \le t \le c = \sqrt{a^2+b^2}\,, \\ 0 & \text{else}\,, \end{cases}$$

where

$$J_{\mathcal{R}}^*(t) = \begin{cases} H_{\mathcal{R},1}^*(t) & \text{if} \quad 0 \le t < a\,, \\ H_{\mathcal{R},2}^*(t) & \text{if} \quad a \le t < b\,, \\ H_{\mathcal{R},3}^*(t) & \text{if} \quad b \le t \le c \end{cases}$$

with

$$\begin{aligned} H_{\mathcal{R},1}^*(t) &= \pi A_{\mathcal{R}} - u_{\mathcal{R}} t + t^2\,, \\ H_{\mathcal{R},2}^*(t) &= -a^2 - 2bt + 2bL_2^*(t,a)\,, \\ H_{\mathcal{R},3}^*(t) &= -(\pi A_{\mathcal{R}} + c^2) - t^2 + 2bL_2^*(t,a) + 2aL_2^*(t,b) \end{aligned}$$

(L_2^* as in our Theorem 2). From this, one easily concludes the distribution function

$$G_{\mathcal{R}}(t) = \int_{-\infty}^{t} g_{\mathcal{R}}(\tau)\,\mathrm{d}\tau = \begin{cases} 0 & \text{if} \quad t < 0\,, \\ \dfrac{2}{A_{\mathcal{R}}^2}\, J_{\mathcal{R}}^\natural(t) & \text{if} \quad 0 \le t < c\,, \\ 1 & \text{if} \quad t \ge c\,, \end{cases}$$

where

$$J_{\mathcal{R}}^\natural(t) = \begin{cases} J_{\mathcal{R},1}^\natural(0,t) & \text{if} \quad 0 \le t < a\,, \\ J_{\mathcal{R},1}^\natural(0,a) + J_{\mathcal{R},2}^\natural(a,t) & \text{if} \quad a \le t < b\,, \\ J_{\mathcal{R},1}^\natural(0,a) + J_{\mathcal{R},2}^\natural(a,b) + J_{\mathcal{R},3}^\natural(b,t) & \text{if} \quad b \le t < c \end{cases}$$

with

$$J^{\natural}_{\mathcal{R},k}(s,t) = H^{\natural}_{\mathcal{R},k}(t) - H^{\natural}_{\mathcal{R},k}(s)\,,$$

where

$$\begin{aligned}
H^{\natural}_{\mathcal{R},1}(t) &= \int t H^{*}_{\mathcal{R},1}(t)\,\mathrm{d}t = \frac{\pi A_{\mathcal{R}} t^2}{2} - \frac{u_{\mathcal{R}} t^3}{3} + \frac{t^4}{4}\,,\\
H^{\natural}_{\mathcal{R},2}(t) &= \int t H^{*}_{\mathcal{R},2}(t)\,\mathrm{d}t = -\frac{a^2 t^2}{2} - \frac{2bt^3}{3} + 2bL^{\natural}_2(t,a)\,,\\
H^{\natural}_{\mathcal{R},3}(t) &= \int t H^{*}_{\mathcal{R},3}(t)\,\mathrm{d}t = -\frac{(\pi A_{\mathcal{R}} + c^2)t^2}{2} - \frac{t^4}{4} + 2bL^{\natural}_2(t,a) + 2aL^{\natural}_2(t,b)
\end{aligned}$$

with $L^{\natural}_2$ as in Corollary 1. Now it is easy to calculate the density function g_2 and the distribution function G_2. Examples are to be found in subsection 5.2.

5 Examples and simulation

5.1 One triangle

Fig. 2 shows some density functions cg. The representation in this diagram has the advantage that cg is independent from the maximum distance c, it depends only on the ratio a/b. Fig. 3 shows the comparison between distribution functions G and corresponding simulation results (plot markers) with $c = 1$ and 10^7 pairs of random points.



Figure 2: $\mathcal{T}_{a,a}$ (thick), $\mathcal{T}_{a,2a}$ (thick and dashed), $\mathcal{T}_{a,5a}$ (thin and dashed) and $\mathcal{T}_{a,20a}$ (thin)



Figure 3: $\mathcal{T}_{a,a}$ (thick) and $\mathcal{T}_{a,5a}$ (thin and dashed)

5.2 Two triangles

Fig. 4 shows the density function cg_2 for $b = 5a$, and the corresponding functions $cg_{\mathcal{R}}$ and cg. Fig. 5 shows the distribution function G_2 for $b = 5a$, and the corresponding functions $G_{\mathcal{R}}$ and G. The plot markers result from a simulation with $c = 1$ and 10^7 pairs of random points.



Figure 4: $cg_{\mathcal{R}}$ (short-dashed), cg (long-dashed) and cg_2 for $b = 5a$



Figure 5: $G_{\mathcal{R}}$ (short-dashed), G (long-dashed) and G_2 for $b = 5a$

References

[1] Ambartzumjan, R. V.; Mecke, J.; Stoyan, D.: *Geometrische Wahrscheinlichkeiten und Stochastische Geometrie,* Akademie Verlag, Berlin, 1993.

[2] Bäsel, U.: Random chords and point distances in regular polygons, arXiv: 1204.2707v2 [math.PR] 5 Sep 2012.

[3] Borel, E.: *Principes et Formules Classiques du Calcul des Probabilités,* Gauthier-Villars, Paris, 1925.

[4] Bronstein, I. N.; Semendjajew, K. A.: *Taschenbuch der Mathematik,* Verlag Nauka, Moskau und BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1989.

[5] Gates, J.: Some properties of chord length distributions, *J. Appl. Prob.,* **24** (1987), 863-874.

[6] Ghosh, B.: Random distances within a rectangle and between two rectangles, *Bull. Calcutta Math. Soc.,* **43** (1951), 17-24.

[7] Gille, W.: The chord length distribution of parallelepipeds with their limiting cases, *Exp. Techn. Phys.,* **36** (1988), 197-208.

[8] Harutyunyan, H. S.; Ohanyan, V. K.: The chord length distribution function for regular polygons, *Adv. Appl. Prob. (SGSA),* **41** (2009), 358-366.

[9] Marsaglia, G.; Narasimhan, B. G.; Zaman, A.: The distance between random points in rectangles, *Commun. Statist. - Theory Meth.,* **19(11)** (1990), 4199-4212.

[10] Mathai, A. M.: *An Introduction to Geometrical Probability,* Gordon and Breach, Australia, 1999.

[11] Mathai, A. M.; Moschopoulos, P.; Pederzoli, G.: Random points associated with rectangles, *Rend. Circ. Mat. Palermo,* Serie II, Tomo XLVIII (1999), 163-190.

[12] Piefke, F.: Beziehungen zwischen der Sehnenlängenverteilung und der Verteilung des Abstandes zweier zufälliger Punkte im Eikörper, *Z. Wahrscheinlichkeitstheorie verw. Gebiete,* **43** (1978), 129-134.

[13] Santaló, L. A.: *Integral Geometry and Geometric Probability,* Addison-Wesley, London, 1976.

[14] Sulanke, R.: Die Verteilung der Sehnenlängen an ebenen und räumlichen Figuren, *Math. Nachr.,* **23** (1961), 51-74.

[15] Zhuang, Y.; Pan, J.: A geometrical probability approach to location-critical network performance metrics, *INFOCOM, Proceedings IEEE,* (2012), 1817-1825.

Uwe Bäsel

HTWK Leipzig, University of Applied Sciences,
Faculty of Mechanical and Energy Engineering,
PF 30 11 66, 04251 Leipzig, Germany,